

MA106 Linear Algebra 2022

Tutorial 3 D1-T4

Kartik Gokhale

Department of Computer Science & Engineering
IIT Bombay

2022

Problem 1

Let A be a $2 \times n$ matrix of real numbers. Is there a relationship between the $\det(AA^T)$ and sum of the (2×2) principal minors of $\det(A^TA)$? Is there a similar such relationship for $3 \times n$ matrices?

Problem 1

Let us brute force it and see what happens

$$A = \begin{bmatrix} u_1 & \cdots & u_n \\ v_1 & \cdots & v_n \end{bmatrix}$$

$$AA^T = \begin{bmatrix} \sum u_i v_i & \sum u_i^2 \\ \sum v_i^2 & \sum u_i v_i \end{bmatrix}$$

$$A^T A = [u_i v_j + u_j v_i]$$

$$M_{kt} = \begin{bmatrix} 2u_k v_k & u_t v_k + v_t u_k \\ u_t v_k + v_t u_k & 2u_t v_t \end{bmatrix}$$

Let us evaluate the sum of determinants

Problem 1

$$\begin{aligned}\sum_{k < t} \det(M_{kt}) &= \sum_{k < t} \det \left(\begin{bmatrix} 2u_k v_k & u_t v_k + v_t u_k \\ u_t v_k + v_t u_k & 2u_t v_t \end{bmatrix} \right) \\ &= \sum_{k < t} u_k v_k u_t v_t - u_t^2 v_t^2 - u_k^2 v_k^2 \\ &= \det(AA^T)\end{aligned}$$

QED

A similar result holds not only for $3 \times n$ matrices but for all $k \times n, k \leq n$ matrices¹

¹Why not for $k > n$?

Problem 2

Show that

$$\det \left(\begin{bmatrix} \cos \alpha & 1 & 0 & \cdots & 0 \\ 1 & 2\cos\alpha & 1 & \cdots & 0 \\ 0 & 1 & 2\cos\alpha & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 2\cos\alpha \end{bmatrix} \right) = \cos n\alpha$$

Problem 2

We will use induction. Trivially, for $n=2$, we have

$$2 \cos^2 \alpha - 1 = \cos 2\alpha$$

By the Induction hypothesis

$$D_k = 2 \cos \alpha \cos((k-1)\alpha) - \cos((k-2)\alpha)$$

Hence,

$$D_k = \cos k \alpha$$

QED

Problem 3

Suppose $\langle x, y \rangle$ is a Hermitian product. Show that

$$4\langle x, y \rangle = \|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2$$

Problem 3

Let us recall some properties about the Hermitian product.

$$\|x\|^2 = \langle x, x \rangle$$

$$\langle x, ay_1 + by_2 \rangle = a\langle x, y_1 \rangle + b\langle x, y_2 \rangle$$

$$\langle ax_1 + bx_2, y \rangle = \bar{a}\langle x_1, y \rangle + \bar{b}\langle x_2, y \rangle$$

Trivial from here

$$\|x + y\|^2 - \|x - y\|^2 = 2\langle x, y \rangle + 2\langle y, x \rangle$$

$$i\|x + iy\|^2 - i\|x - iy\|^2 = 2\langle x, y \rangle - 2\langle y, x \rangle$$

$$4\langle x, y \rangle = \|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2$$

QED

Problem 4

Sheet 4 Problem 4

Problem 4

$$\{[1, 1, 0, 0], \frac{1}{2}[1, -1, 2, 0], \frac{1}{3}[1, -1, -1, 3], \frac{1}{2}[-1, 1, 1, 1], \mathbf{0}, \mathbf{0}\}.$$

The 4 nonzero elements give an orthogonal basis. $[-2, -1, 1, 2]$ must be in linear span.

To verify Bessel's equality, we first note that

$$\mathbf{u}_1 = \frac{1}{\sqrt{2}}[1, 1, 0, 0], \quad \mathbf{u}_2 = \frac{1}{\sqrt{6}}[1, -1, 2, 0], \quad \mathbf{u}_3 = \frac{1}{2\sqrt{3}}[1, -1, -1, 3], \quad \mathbf{u}_4 = \frac{1}{2}[-1, 1, 1, 1].$$

$$\text{Now } \sum (\mathbf{v} \cdot \mathbf{u}_j)^2 = \frac{9}{2} + \frac{1}{6} + \frac{4}{3} + 4 = 10 = \|\mathbf{v}\|^2. \text{ (Verified!)}$$

Problem 5

Sheet 4 Problem 8

Problem 5

$$\begin{aligned}\mathbf{w}_1 = \mathbf{v}_1 &= [1, i, 0, 0, 0] \\ \mathbf{w}_2 = [0, 1, i, 0, 0] - \frac{-i}{2}[1, i, 0, 0, 0] &= \frac{1}{2}[i, 1, 2i, 0, 0] \\ \text{Similarly, } \mathbf{w}_3 &= \frac{1}{3}[-1, i, 1, 3i, 0] \\ \text{and } \mathbf{w}_4 &= \frac{1}{4}[-i, -1, i, 1, 4i].\end{aligned}$$

The corresponding orthonormal (unitary) set is

$$\left\{ \frac{1}{\sqrt{2}}[1, i, 0, 0, 0], \frac{1}{\sqrt{6}}[i, 1, 2i, 0, 0], \frac{1}{2\sqrt{3}}[-1, i, 1, 3i, 0], \frac{1}{2\sqrt{5}}[-i, -1, i, 1, 4i] \right\}.$$

$$\|[1, i, 1, i, 1]\|^2 = 5. \text{ OTOH,}$$

$$|\langle \mathbf{v}, \mathbf{u}_1 \rangle|^2 = 2, \quad |\langle \mathbf{v}, \mathbf{u}_2 \rangle|^2 = 2/3,$$

$$|\langle \mathbf{v}, \mathbf{u}_3 \rangle|^2 = 4/3, \quad |\langle \mathbf{v}, \mathbf{u}_4 \rangle|^2 = 4/5.$$

The sum is $24/5 < 5$, hence can not be in the linear span.

Problem 6

Show that if A is $n \times n$ complex matrix whose rows are orthonormal, then so are its columns

Problem 6

For the beauty of this concise proof, let me represent OC as orthonormal columns and OR as orthonormal rows

$$\text{OR} \implies (AA^\dagger = I) \implies (A^\dagger = A^\dagger AA^\dagger) \implies (A^\dagger A = I) \implies \text{OC}$$

Problem 7

$$V = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ i \\ -1 \end{bmatrix}, W = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ -2i \\ -1 \end{bmatrix}$$

Find $U \in \mathcal{C}^3$ such that the 3 are orthonormal. Any other way than Gram Schmidt?

Problem 7

A trivial method exists using Gram Schmidt Orthonormalisation. Consider the trivial basis of \mathcal{C}^3 and orthormalize them with respect to the given vectors. At least one is bound to give you U
Another method would be to use the results of Problem 6.
Construct a vector U such that the augmented matrix formed has orthonormal rows.

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Can you think of any other method?